



IRN France-Japan Frontiers in Plant Biology

IRN-FJFPB

Webinar « Development and Adaptation »

October 21st, 2021

Caroline Bellenot (PhD student), LIPME, Université de Toulouse, INRAE/CNRS, UMR 0441/2598, F-31326 Castanet-Tolosan, France

Contact laurent.noel@inrae.fr

<http://www.xantho.fr>

Caroline Bellenot¹, Alexandra Guowietz², Gilles Clément³, Sylvie Citerne³, Magalie Charvin⁵, Marie-Françoise Jardinaud¹, Sébastien Carrère¹, Nathalie Leonhardt⁴, Laurent Nussaume⁴, Grégory Mouille³, Lionel Navarro⁵, Martin Schattat², Jean-Marc Routaboul^{1*}, Laurent, D. Noël¹

Affiliations

¹ LIPME, Université de Toulouse, INRAE/CNRS, UMR 0441/2598, F-31326 Castanet-Tolosan, France
LBDP

² Department of Plant Physiology, Institute for Biology, Martin-Luther-University Halle-Wittenberg, D-06120 Halle (Saale), Germany

³ Institut Jean-Pierre Bourgin, INRA and AgroParisTech and CNRS, Université Paris-Saclay, F-78000 Versailles, France

⁴ Laboratoire de Biologie du Développement des Plantes, Institut de Biosciences et

Biotechnologies d'Aix-Marseille, Aix-Marseille Université and Commissariat à l'Energie Atomique et aux Énergies Alternatives and CNRS, UMR 7265, F-13108 Saint Paul-Les-Durance, France

⁵ Institut de Biologie de l'Ecole Normale Supérieure (IBENS), Centre National de la Recherche Scientifique UMR8197, Institut National de la Santé et de la Recherche Médicale U1024, 75005 Paris, France.

Physiology and immunity of *Brassicaceae* hydathodes upon infection by bacterial pathogens

Hydathodes are leaf organs present in all vascular plants where they mediate guttation. These organs provide a facilitated access to plant inner tissues including xylem vessels. Hydathodes are thus preferred infection sites for several bacterial vascular pathogens such as *Xanthomonas campestris* pv. *campestris* (*Xcc*), the causal agent of black rot disease in *Brassicaceae*. Our goal is to unravel the physiology and immunity of hydathodes in *Arabidopsis* and cauliflower using metabolomic and transcriptomic approaches in healthy and infected hydathodes. Latest results will be presented.



INRAE



Université Claude Bernard Lyon 1

